


$\sigma$  and  $R$  in  $e^+e^-$  Collisions

**Figure 39.6, Figure 39.7:** World data on the total cross section of  $e^+e^- \rightarrow \text{hadrons}$  and the ratio  $R = \sigma(e^+e^- \rightarrow \text{hadrons})/\sigma(e^+e^- \rightarrow \mu^+\mu^-)$ , QED simple pole). The curves are an educative guide. The solid curves are the 3-loop pQCD predictions for  $\sigma(e^+e^- \rightarrow \text{hadrons})$  and the  $R$  ratio, respectively [see our Review on Quantum chromodynamics, Eq. (9.12)] or, for more details, K.G. Chetyrkin *et al.*, Nucl. Phys. **B586**, 56 (2000), Eqs. (1)–(3). Breit-Wigner parameterizations of  $J/\psi$ ,  $\psi(2S)$ , and  $\Upsilon(nS)$ ,  $n = 1..4$  are also shown. **Note:** The experimental shapes of these resonances are dominated by the machine energy spread and are not shown. The dashed curves are the naive quark parton model predictions for  $\sigma$  and  $R$ . The full list of references, as well as the details of  $R$  ratio extraction from the original data, can be found in O.V. Zenin *et al.*, hep-ph/0110176 (to be published in J. Phys. **G**). Corresponding computer-readable data files are available at [http://wwwppds.ihep.su/~zenin\\_o/contents\\_plots.html](http://wwwppds.ihep.su/~zenin_o/contents_plots.html). (Courtesy of the COMPAS (Protvino) and HEPDATA (Durham) Groups, November 2001.)